



**Acta Botanica  
Mexicana**

# Mycelial inhibition of *Trichoderma* spp. (Hypocreaceae) isolated from the cultivation of *Pleurotus ostreatus* (Pleurotaceae) with an extract of *Pycnoporus* sp. (Polyporaceae)

## Inhibición micelial de *Trichoderma* spp. (Hypocreaceae) aislado del cultivo de *Pleurotus ostreatus* (Pleurotaceae) con un extracto de *Pycnoporus* sp. (Polyporaceae)

Anaid Talavera-Ortiz<sup>1,2</sup> , Priscila Chaverri<sup>3</sup> , Gerardo Díaz-Godínez<sup>4</sup>   
Ma. de Lourdes Acosta-Urdapilleta<sup>1</sup> , Elba Villegas<sup>5</sup> , Maura Téllez-Téllez<sup>1,6</sup>

### Abstract:

**Background and Aims:** The production of edible fungi is affected by bacterial, fungal and viral diseases, which very often cause large losses. In the production of mushrooms of the genus *Pleurotus*, the fungi of *Trichoderma* spp. represent a serious problem of contamination and although there are some chemical compounds that control the infection, they are not entirely safe for human consumption. As a consequence, alternatives are being searched for through biotechnology, such as the one presented in this paper.

**Methods:** Strains of fungi of the genus *Trichoderma* were isolated from the substrate where *Pleurotus ostreatus* was being cultivated. These were identified morphologically and molecularly, followed by tests to inhibit the growth of *Trichoderma* strains in both agar and wheat straw, using a cetonc extract of the dehydrated fruiting body of *Pycnoporus* sp.

**Key results:** Two strains of *Trichoderma* (*T. pleuroti* and *T. atroviride*, belonging to the clade of *T. harzianum*) were isolated from infected substrate obtained in production modules of *Pleurotus ostreatus* located in Tlaquitenango and Cuernavaca, in the state of Morelos, Mexico. The effect of a cetonc extract of the fruiting body of *Pycnoporus* sp. on the mycelial growth of the isolated strains of *Trichoderma* was also evaluated, observing decrease in mycelial growth rate in Petri dish up to 72% and on lignocellulosic substrate both mycelial growth and sporulation were delayed up to 10 days.

**Conclusions:** The extract of *Pycnoporus* sp. could be an alternative to control the infection by *Trichoderma* spp. in mushroom cultures of the genus *Pleurotus*.

**Key words:** disease control, edible mushrooms, lignocellulosic substrate.

### Resumen:

**Antecedentes y Objetivos:** La producción de hongos comestibles se ve afectada por enfermedades bacterianas, fúngicas y virales, que a menudo causan grandes pérdidas. En la producción de hongos del género *Pleurotus*, los hongos de *Trichoderma* spp. representan un grave problema de contaminación y, aunque hay algunos compuestos químicos que controlan la infección, no son completamente seguros para el consumo humano. Como consecuencia, se están buscando alternativas a través de la biotecnología, como la que se presenta en este manuscrito.

**Métodos:** Cepas de hongos del género *Trichoderma* se aislaron del sustrato donde se cultivaba *Pleurotus ostreatus*. Estos se identificaron morfológicamente y molecularmente; posteriormente se hicieron pruebas para inhibir el crecimiento de las cepas de *Trichoderma* tanto en agar como en paja de trigo, utilizando un extracto cetónico del cuerpo fructífero deshidratado de *Pycnoporus* sp.

**Resultados clave:** Se aislaron dos cepas de *Trichoderma* (*T. pleuroti* y *T. atroviride*, perteneciente al clado de *T. harzianum*) del sustrato infectado obtenido en los módulos de producción de *Pleurotus ostreatus* ubicados en Tlaquitenango y Cuernavaca, en el estado de Morelos, México. También se evaluó el efecto de un extracto cetónico del cuerpo fructífero de *Pycnoporus* sp. En el crecimiento micelial de las cepas aisladas de *Trichoderma*, observando una disminución en la tasa de crecimiento del micelio en la placa de Petri hasta en 72%, y en el sustrato lignocelulósico, tanto el crecimiento micelial como la esporulación se retrasaron hasta 10 días.

**Conclusiones:** El extracto de *Pycnoporus* sp. podría ser una alternativa para controlar la infección por *Trichoderma* spp. en cultivos de hongos del género *Pleurotus*.

**Palabras clave:** control de enfermedades, hongos comestibles, sustrato lignocelulósico.

<sup>1</sup> Universidad Autónoma del Estado de Morelos, Centro de Investigaciones Biológicas, Av. Universidad No. 1001, Col. Chamilpa, 62209 Cuernavaca, Morelos, Mexico.

<sup>2</sup> Universidad Autónoma del Estado de Morelos, Centro de Investigaciones Biológicas, Maestría en Manejo de Recursos Naturales, Av. Universidad No. 1001, Col. Chamilpa, 62209 Cuernavaca, Morelos, Mexico.

<sup>3</sup> Universidad de Costa Rica, Escuela de Biología, Ciudad Universitaria Rodrigo Facio Brenes, San Pedro de Montes de Oca, San José, Costa Rica.

<sup>4</sup> Universidad Autónoma de Tlaxcala, Centro de Investigación en Ciencias Biológicas, Mariano Matamoros Periferia, Villa Mariano Matamoros, 90120 Tlaxcala, Tlaxcala, Mexico.

<sup>5</sup> Universidad Autónoma del Estado de Morelos, Centro de Investigación en Biotecnología, Av. Universidad No. 1001, Col. Chamilpa, 62209 Cuernavaca, Morelos, Mexico.

<sup>6</sup> Author for correspondence: [maura.tellez@uaem.mx](mailto:maura.tellez@uaem.mx)

Received: March 31, 2019.

Reviewed: May 28, 2019.

Accepted by Marie-Stéphanie Samain: July 29, 2019.

Published Online first: August 15, 2019.

Published: Acta Botanica Mexicana 127 (2020).

To cite as:

Talavera-Ortiz, A., P. Chaverri, G. Díaz-Godínez, M. L. Acosta-Urdapilleta, E. Villegas and M. Téllez-Téllez. 2019(2020). Mycelial inhibition of *Trichoderma* spp. (Hypocreaceae) isolated from the cultivation of *Pleurotus ostreatus* (Pleurotaceae) with an extract of *Pycnoporus* sp. (Polyporaceae). Acta Botanica Mexicana 127: e1537. DOI: [10.21829/abm127.2020.1537](https://doi.org/10.21829/abm127.2020.1537)



This is an open access article under the Creative Commons 4.0 Attribution-Non commercial Licence (CC BY-NC 4.0 International).

e-ISSN: 2448-7589

## Introduction

Among the edible fungi cultivated worldwide with higher production are *Agaricus bisporus* (J.E. Lange) Imbach (Agaricaceae) and *Pleurotus ostreatus* (Jacq.) P. Kumm. (Pleurotaceae; both families belonging to Agaricales, Agaricomycetidae, Agaricomycetes, Agaricomycotina, Basidiomycota, Fungi; [Index Fungorum](#), 2019) ([Royse](#), 2014), which present diseases and pests such as green mold caused by species of *Trichoderma* Pers. (Hypocreaceae) ([Oei](#), 1991; [Park et al.](#), 2004), fire mold by *Neurospora* sp. (Sordariaceae) and brown spot of *Pseudomonas tolaassi* Paine, as well as some by viruses ([Kim et al.](#), 1995; 2000).

The disease generated by fungi of the genus *Trichoderma* is difficult to distinguish, because in the culture of mushrooms of the genus *Pleurotus* (Fr.) P. Kumm. (Pleurotaceae), both mycelia have similar characteristics, so the infection is detectable after approximately two weeks only, as it turns green due to the strong sporulation ([Danesh et al.](#), 2000), causing the inhibition of mycelial growth of *Pleurotus* because they compete for space and nutrients. The strategy of *Trichoderma* species is the secretion of hydrolytic enzymes, such as chitinases,  $\beta$ -glucanases and cellulases, which lyse the cell walls of the *Pleurotus* mycelium ([Goltapeh and Danesh](#), 2000), causing the fruiting bodies not to form ([Beyer et al.](#), 2000).

The first report of *Trichoderma* infection in *P. ostreatus* culture occurred in North America ([Sharma and Vijay](#), 1996), then in Italy ([Woo et al.](#), 2004), South Korea ([Park et al.](#), 2006), Romania ([Kredics et al.](#), 2006), Hungary ([Hatvani et al.](#), 2007) and Spain ([Gea](#), 2009). The species of *Trichoderma* are asexual and they inhabit the soil ([Kredics et al.](#), 2009). The main *Trichoderma* species that cause green mold disease in mushroom cultures of the genus *Pleurotus* are *T. pleuroticola* S.H. Yu & M.S. Park ([Hatvani et al.](#), 2007), *T. harzianum* Rifai ([Sobieralski et al.](#), 2012) and *T. pleuroti* S.H. Yu & M.S. Park (as 'pleurotum') (Hypocreaceae, Hypocreales, Hypocreomycetidae, Sordariomycetes, Pezizomycotina, Ascomycota, Fungi; [Index Fungorum](#), 2019) ([Marik et al.](#), 2017). It has been estimated that *Trichoderma*, together with mushroom virus X (MVX is a relatively new disease that affects the production of the commercial mushrooms), has caused a 25% loss in the production of cultivated edible mushrooms ([Soković and van Griensven](#), 2006; [Sobieralski et al.](#), 2012).

*Trichoderma pleuroti* and *T. pleuroticola* have been found in the substrate of mushroom cultivation of the genus *Pleurotus* in different countries; however, the second species has also been found in soil and wood, as well as associated with *P. ostreatus* that grows in natural environments in countries such as Canada, United States of America, Iran, New Zealand and some European ones ([Komoń-Zelazowska et al.](#), 2007; [Kredics et al.](#), 2009). This represents a higher risk of infection for the mushroom cultivation, since the sources of infection by *T. pleuroticola* can be diverse ([Kredics et al.](#), 2009). *Trichoderma pleuroti* could be specialized to an ecological niche related to substrates of mushroom cultivation of the genus *Pleurotus* ([Komoń-Zelazowska et al.](#), 2007). One way to control the disease caused by *Trichoderma* has been to use disinfectants (chlorine, calcium hydroxide) and/or fungicides; the latter could generate resistant strains that require more of these chemical compounds to control the problem. In this sense it is very important to look for alternatives of biological control of the disease of the green mold, knowing the *Trichoderma* species involved in the cultures of *P. ostreatus*.

It is worth mentioning that the fungi of the genus *Pycnoporus* P. Karst. (Polyporaceae, Polyporales, Incertae sedis, Agaricomycetes, Agaricomycotina, Basidiomycota, Fungi; [Index Fungorum](#), 2019), although not considered edible due to their corky texture, have medicinal and biological properties, such as antimicrobial activity according to reports in China, Japan and the rest of Asia (although less frequent) ([Calonge](#), 2011). In Mexico it has been used to combat acne, warts, diarrhea, dysentery and tinea capitis (cutaneous mycosis) ([Martínez-Alfaro et al.](#), 1983; [Guzmán](#) 1994; 2003; 2008). Currently, there are no reports on the use of *Pycnoporus* species against fungi of the genus *Trichoderma*, but there are a few that indicate activity against other organisms of the same kingdom. Recently, it was informed that the culture broth of *P. cinnabarinus* (Jacq.) P. Karst. had a bacteriostatic and bactericidal effect against *Escherichia coli* Escherich and *Staphylococcus aureus* Rosenbach, respectively, suggesting that cinnabarin and cinnabarinic acid are the compounds produced by the fungus with antimicrobial activity ([Díaz-Godínez et al.](#), 2016). [Al-Fatimi et al.](#) (2013) reported that the methanolic extract of fruiting bodies of *Pycnoporus sanguineus* (L.) Murrillo showed con-

siderable antifungal activity against the human pathogens *Candida albicans* (C.P. Robin) Berkhout, *C. krusei* (Castell.) Berkhout, *Microsporium gypseum* (E. Bodin) Guiart & Grigoraki, *Trichophyton mentagrophytes* (C.P. Robin) Sabour. and against *C. maltosa* Komag., Nakase & Katsuya, which is not a pathogenic fungus for humans. In another study, aqueous and methanolic extracts of the mycelium of *P. sanguineus* showed antifungal activity against different species of wood-degrading fungi; the methanolic extract had greater inhibition, particularly against *Lentinus* sp., *Microporus affinis* (Blume & T. Nees) Kuntze, *M. xanthopus* (Fr.) Kuntze and *Trametes menziesii* (Berk.) Ryvarden, but presented a lesser effect against *Schizophyllum commune* Fr. (Teoh, 2011). Based on the above, it was proposed to use extracts of fruiting bodies of *Pycnoporus* sp. to try to inhibit the growth of *Trichoderma* strains isolated from cultures of *Pleurotus ostreatus*.

## Materials and Methods

### Isolation of *Trichoderma* strains

Isolates of *Trichoderma* spp. were obtained by collecting spores of the mycelium that showed typical characteristics of the disease of green mold on the substrate of the culture of *Pleurotus ostreatus*. Sampling was carried out in eight production modules located in the municipalities of Tlaquitenango and Cuernavaca, Morelos, Mexico. Dextrose potato agar (PDA, Difco Laboratories) was used at the beginning of the isolation process of the strains of *Trichoderma* spp. (Chen et al., 1999). The inoculated plates were incubated at 25 °C (Incubator, LabTech LIB-150M, Jalisco, Mexico) in the dark, until mycelial growth was observed. To assure the process, the hyphal tip isolation was carried out, which consisted in taking with a sterile needle a portion of the sporulated mycelium, followed by inoculation in Petri dishes with water-agar, and incubation. After 24 h, the tip of a hypha was cut with a dissecting needle under a stereoscopic microscope (Motic SMZ 161, Mexico City, Mexico), placed on PDA, and then incubated at 25 °C in darkness (Zehr, 1978).

### Molecular identification and morphological characteristics of *Trichoderma* strains

Mycelia were obtained by growing each *Trichoderma* strain isolated in potato dextrose broth by means of the in-

clined tube technique. For DNA extraction from previously dried mycelium, the PrepMan® Ultra Sample Preparation Reagent KIT (Thermo Fisher Scientific, Waltham, USA) was used following the manufacturer's instructions. The Translation Elongation Factor 1-alpha was amplified by PCR using the primers Tef 728M-f (5'-CAT YGA GAA GTT CGA GAA GG-3') and Ef2r (5'-GGA RGT ACC AGT SAT CAT GTT-3'), which give a product of 400 bp. The 50 µl reaction mixture contained 1 pmole of both primers, 2.0 U of Taq DNA polymerase (Promega), 0.1 µl of 4 mM dNTP, 5 µl of 10X buffer and 100 ng of DNA. The parameters for PCR amplification were 94 °C for 2 min; 35 cycles of 94 °C for 40 s, 66 °C for 55 s, 72 °C for 2 min and final extension 72 °C for 8 min, were cleaned with ExoSAP-IT (USB Corp., Cleveland, USA). The sequencing was performed in Macrogen, Maryland, USA. The rDNA sequences were compared with the data already published in GenBank database using the tool BLAST (NCBI, 2019). The morphological characteristics of each of the strains were also determined using an optical microscope (Motic BA310, Mexico City, Mexico) by means of preparations with 3% KOH, and conidia, phialides and conidiophores were measured. The keys used for their identification were those described by Samuels and Prakash (2015).

### Mycelial growth rate of *Trichoderma* strains

The mycelial growth rate (Vc) of two strains of *Trichoderma* isolated (Tr1 and Tr5, each one as representative of the two determined species, see Table 1), as well as of *T. aggressivum* Samuels & W. Gams (Hypocreaceae, Hypocreales, Hypocreomycetidae, Sordariomycetes, Pezizomycotina, Ascomycota, Fungi; Index Fungorum, 2019) (CPM-113, donated by CREGENHC (Centro de Recursos Genéticos de Hongos Comestibles del Colegio de Postgraduados), which was used as a positive control) in PDA, with and without photoperiod was measured. Each of the strains was inoculated (4 mm plug) in Petri dishes with PDA, incubated at 25 °C. One experiment was performed in complete darkness and another with photoperiod (12 h light/12 h dark). Each experiment was performed in triplicate and the mean ± SD of the Vc was reported in mm/h (Téllez-Téllez et al., 2003). An ANOVA and Tukey test using STATISTICA V. 7.0 (StatSoft Inc., 2004) were also performed.

**Table 1:** Molecular identification and morphological characteristics of strains of *Trichoderma* Pers. (Hypocreaceae) isolated from the culture substrate of *Pleurotus ostreatus* (Jacq.) P. Kumm.

Molecular identification				Morphological characteristics
Strain	Identity (%) GenBank	NCBI GenBank accession number	Species	The values are long × wide (minor value) average (higher value)
Tr1	97	HM142382.1	<i>T. pleuroti</i> S.H. Yu & M.S. Park	Green color <i>Gliocladium</i> Corda type conidiophore Conidia ellipsoidal shape, (2.9-)3.0-3.2(-3.6) × (1.9-)2.0-2.1(-2.5) μm Phialides, (11.5-)11.9-12.0(-12.2) × (2.5-)2.6-2.7(-2.8), in 45° position of the base of the conidiophore
Tr2	95	HM142382.1	<i>T. cf. pleuroti</i> S.H. Yu & M.S. Park	Green color <i>Gliocladium</i> Corda type conidiophore Conidia ellipsoidal shape, (2.9-)3.0-3.1(-3.3) × (1.9-)2.0-2.2(-2.5) μm Phialides, (7.0-)7.5-7.8(-8.0) × (1.9-)2.0-2.7(-3.0) μm
Tr3	99	KJ665413.1	<i>T. atrobrunneum</i> F.B. Rocha, P. Chaverri & Jaklitsch	Green color Pyramidal type conidiophore Conidia subglobose shape, (2.8-)3.2-3.5(-4.0) × (2.5-)2.8-3.0(-3.6) μm Phialides, (5.0-)5.2-5.5(-6.0) × (2.6-)2.9-3.0(-3.2) μm
Tr4	99	KJ665413.1	<i>T. atrobrunneum</i> F.B. Rocha, P. Chaverri & Jaklitsch	Green color Pyramidal type conidiophore Conidia subglobose shape, (2.8-)3.2-3.5(-4.0) × (2.5-)2.8-3.0(-3.6) μm Phialides, (5.0-)5.2-5.5(-6.0) × (2.6-)2.9-3.0(-3.2) μm
Tr5	99	KJ665413.1	<i>T. atrobrunneum</i> F.B. Rocha, P. Chaverri & Jaklitsch	Green color Pyramidal type conidiophore Conidia subglobose shape, (2.8-)3.2-3.5(-4.0) × (2.5-)2.8-3.0(-3.6) μm Phialides, (5.0-)5.2-5.5(-6.0) × (2.6-)2.9-3.0(-3.2) μm
Tr6	96	HM142382.1	<i>T. cf. pleuroti</i> S.H. Yu & M.S. Park	Green color <i>Gliocladium</i> Corda type conidiophore Conidia: ellipsoidal shape, (2.3-)2.5-3.0(-3.3) × (1.5-)1.8-2.0(-2.4) μm Phialides, (5.5-)6.0-6.8(-7.0) × (2.7-)3.0-3.5(-3.8) μm

### Inhibition of the growth of *Trichoderma* strains

An extract with acetone was obtained from the dry fruiting body of *Pycnoporus* sp. (strain HEMIM-51 donated by Herbario Micológico de Morelos); the mixture was stirred at 125 rpm for 12 h, then the solvent was removed by a rotary evaporator (Janke & Kunkel GMBH-IKA Labortechnik RV 06 System, Staufen, Germany). Inhibition tests were performed in both Petri dishes with agar and on lignocellulosic substrate. In the first case, different concentrations of the extract (1.2, 2.4 and 3.6% v/v) were added to the PDA and the strains isolated from *Trichoderma* (Tr1 and Tr5) and *T. aggressivum* were inoculated separately. The strain of *P. ostreatus* (HEMIM-50) was also used to determine the effect

of the extract on its growth. All the cultures were incubated at 25 °C and the Vc was measured every 12 h until total invasion of the plate. The percentage of inhibition was calculated with the following formula:

$$\% \text{ inhibition} = \frac{V - V_e \times 100}{V}$$

Where:

V: Vc without extract

Ve: Vc with extract

For tests on lignocellulosic substrate, 10 g of wheat straw (75% humidity) was placed in cylindrical glass con-

tainers with a nominal volume of 250 ml, then sterilized (90 min/120 °C). To each experimental unit the extract of *Pycnoporus* sp. was added at 3.6% (v/w) considering the weight of straw. A parallel experiment was carried out without the addition of the extract as control. Consequently, the *Trichoderma* strains isolated (Tr1 and Tr5) were inoculated separately and incubated at 25 °C during 10 days. Mycelial invasion and sporulation were observed in comparison with the control.

## Results

In the different production modules of *P. ostreatus*, the characteristic symptoms of *Trichoderma* disease were observed in the substrates. Initially, there was presence of compact white mycelium that is confused with the *Pleurotus* mycelium and which gradually turns green, due to the sporulation. The isolation of *Trichoderma* strains was achieved by obtaining spores only in six of the eight samples. The six *Trichoderma* strains isolated were called Tr1, Tr2, Tr3, Tr4, Tr5 and Tr6. Through the molecular identification, the strains Tr1, Tr2 and Tr6 showed an identity of 97, 95 and 96%, respectively, with *T. pleuroti*, reason why the first strain was considered as *T. pleuroti* and the last two as *T. cf. pleuroti*. Three strains showed an identity of 99% with *T. atrobrunneum* F.B. Rocha, P. Chaverri & Jaklitsch (Hypocreaceae, Hypocreales, Hypocreomycetidae, Sordariomycetes, Pezizomycotina, Ascomycota, Fungi; [Index Fungorum, 2019](#)) (Tr3, Tr4 and Tr5). In general, the strains presented typical morphological characteristics of each species ([Table 1](#)).

It was decided to work with Tr1 and Tr5, since the first presented the highest percentage of similarity with *T. pleuroti*, and the second was used as a representative strain of those identified as *T. atrobrunneum*, considering the same strain as Tr3 and Tr4. The culture in photoperiod or in darkness caused changes in the pigmentation of the mycelium. In general those with photoperiod showed a distinctive yellow coloration of young conidia; however, in the dark, they presented the typical green pigmentation ([Fig. 1](#)). The Vc of Tr1 and Tr5 were 0.52 and 0.55 mm/h in photoperiod, respectively, and in darkness 0.32 and 40 mm/h, for the same strains. For *T. aggressivum* with photoperiod Vc was 0.57 and in the dark 0.49 mm/h.


[Figure 2](#) shows the inhibition of mycelial growth of *Trichoderma* and *P. ostreatus* assessed in Petri dishes with the extract of *Pycnoporus* sp. A clear effect of the extract on the *Trichoderma* strains was observed, while the mycelial growth of *Pleurotus* was minimally affected (8%) by the highest concentration of the extract, and no effect was observed with the other two concentrations.

In general, the effect of the extract on *Trichoderma* strains was concentration dependent; Tr5 and *T. aggressivum* were the most affected, with 1.2% of the extract retarded mycelial growth in 48 and 33%, respectively. With concentrations of 2.4 and 3.6% the inhibition was 61-74% without significant difference ( $p < 0.05$ ). Strain Tr1 was also affected by the extract, observing inhibition of 21, 27 and 38% with concentrations of 1.2, 2.4 and 3.6%, respectively.


Strains Tr1 and Tr5 grown in wheat straw had abundant mycelium after 10 days of culture ([Figs. 3A, C](#)), while the addition of the extract of *Pycnoporus* sp. (3.6 %) completely inhibited their growth ([Figs. 3B, D](#)). The strain of *T. aggressivum* was less affected by the extract; at 10 days of incubation, the sporulation was delayed by the extract, since the typical green pigmentation could be observed in the absence of the extract ([Fig. 3E](#)) and only mycelium in the presence of the extract ([Fig. 3F](#)). *Pleurotus ostreatus* showed abundant mycelium ([Fig. 3G](#)); however, when the extract was added, the mycelial invasion was delayed by ten days ([Fig. 3H](#)).

## Discussion

The disease caused by *Trichoderma* strains has been reported in several parts of the world, among them Europe, Asia, North America and Latin America; many studies have been realized to identify the species causing this problem, specifically for the cultivation of *P. ostreatus*. [Choi et al. \(2003\)](#) reported isolates of *Trichoderma* spp. obtained from *P. ostreatus* and *P. eryngii* (DC.) Qué. cultures using rice straw, cotton waste and sawdust as substrates. The strains found were *Trichoderma cf. virens* (J.H. Mill., Giddens & A.A. Foster) Arx (70.8%), *T. longibrachiatum* Rifai (16.7%) and *T. harzianum* (12.5%). [Park et al. \(2004\)](#) performed the molecular and morphological analyses of two isolates of *Trichoderma* from cultures of oyster mushroom (*P. ostreatus*) in Korea. These were proposed as new species with the



**Figure 1:** Mycelial growth of strains of *Trichoderma* Pers. (Hypocreaceae) in PDA. A. Tr1 (darkness); B. Tr1 (photoperiod); C. Tr5 (darkness); D. Tr5 (photoperiod) and *Trichoderma aggressivum* Samuels & W. Gams in E. darkness and in F. photoperiod.



**Figure 2:** Inhibition of mycelial growth of strains of *Trichoderma* Pers. (Hypocreaceae) and *Pleurotus ostreatus* (Jacq.) P. Kumm. on agar in Petri dishes added with the extract of *Pycnoporus* sp.

names *T. koreanum* S.-Y. Oh, M.S. Park & Y.W. Lim and *T. pleuroti*, K1 and K2, respectively. Later, Park et al. (2006) renamed these two species as *T. pleurotum* and *T. pleuroticola*. It should be mentioned that *T. pleuroticola* is associated with the cultures of all mushrooms of the genus *Pleurotus*, while *T. pleurotum* has been found only in *P. ostreatus* cultures, both wild and cultivated, as well as in soil and wood in countries such as Canada, United States of America, Iran, New Zealand and some European ones (Kredics et al., 2009). Komoń-Zelazowska et al. (2007) informed that the origin of infection by *T. pleurotum* (*T. pleuroti*) was through the substrate, so it must have an adequate management of straws used in the cultivation of mushrooms to reduce the frequency of the appearance of the disease. *Trichoderma harzianum* has been observed on decaying wood and associated with mushrooms in North America and Europe (Chaverri et al., 2015) and *T. aggressivum* in mushroom cultivation in Europe, Iran and South Korea (Komoń-Zelazowska et al., 2007).

Sobal et al. (2016) conducted a study in Mexico where they isolated 24 strains of *Trichoderma* from substrates of commercial cultivation of edible mushrooms in the states of Puebla, Tlaxcala, Morelos, State of Mexico and Veracruz, identifying *T. aggressivum* f. *agressivum*, *T. koningii* Oudem, *T. harzianum*, *T. atroviride* Bissett, *T. citrinoviride* Bissett and *T. pleuroti*. In our work, *T. pleuroti* and *T. atroviride* were found in the state of Morelos, and it is important to mention

that the latter was not reported by Sobal et al. (2016), so this contributes to the knowledge of the species that cause diseases in the cultivation of mushrooms of the genus *Pleurotus* in the central region of Mexico.

It was communicated that *T. pleuroticola* showed a Vc similar to that of *T. aggressivum* while *T. pleurotum* was slower (Komoń-Zelazowska et al., 2007). In this work, it was observed that in general *T. aggressivum* presented higher Vc compared with *T. pleuroti*. Chaverri et al. (2015) documented that *T. atroviride* grown on PDA after 96 h at 25 °C, presented abundant aerial and cottony or woolly mycelium, the conidia were abundant within 48 to 72 h in wide concentric rings, which coincides with the characteristics presented by the Tr5 strain.

Qiu et al. (2017) said that the high temperature favored a greater production of conidia in less time for *T. asperellum* Samuels, Lieckf. & Nirenberg, together with the hyphae of *P. ostreatus*. In this work three strains of *T. atroviride* were isolated; Tr3 and Tr4 from the northern area of the municipality of Cuernavaca, Morelos, where the temperature ranges from 15-31 °C, and Tr5 from the municipality of Tlaquitenango, where the temperature is 19-35 °C. It is suggested that the Tr5 strain grew faster and sporulated (presented the green coloration) in less time, in comparison with Tr3 and Tr4 (data not shown), given the temperature of the place of origin that reaches up to 35 °C, although more tests that can confirm the effect of temperature should be performed.

Given that the disease of green mold causes great losses in the production of mushrooms of the genus *Pleurotus*, many works have been carried out with the purpose of establishing strategies for its control, including the use of commercial fungicides and natural compounds, as well as the modification of the temperature and the pH of the cultures. However, this has had little success since it affects both the guest and the host (Woo et al., 2004; Wang et al., 2016). Hatvani et al. (2012) reported that certain natural compounds of phenolic type such as thymol, ferulic acid, (+)-menthol, and (-)-menthol had an inhibitory effect on the growth of *T. pleurotum*, *T. pleuroticola* and *T. harzianum*; however, these also inhibited the growth of the edible mushrooms.

Reyes (2013) tested an ethanolic extract of *Ruta graveolens* L. (4 g/L) in *T. aggressivum* (CPM-113), obtain-



**Figure 3:** Growth of strains Tr1, Tr5, *Trichoderma aggressivum* Samuels & W. Gams and *Pleurotus ostreatus* (Jacq.) P. Kumm, without (A, C, E and G, respectively) and with (B, D, F and H, respectively) the addition of the extract of *Pycnoporus* sp. (Polyporaceae).

ing an inhibition of 53% at 24 h, and 44% after 48 h, and they found that the fungus adapted to the presence of the compound over time, with abundant mycelium and change in the color of the spores. This phenomenon was not observed in any of the strains of *Trichoderma* studied in this work; only delays were obtained in the mycelial growth rate and as a consequence also in sporulation, the latter happened both on plate and on wheat straw. After ten days of incubation on wheat straw, the strain *T. aggressivum* (CPM-113) did not show the green coloration with the extract, while without the addition of the extract the sporulation appeared, which indicates that although it does not completely inhibit this strain, its growth and sporulation were slower. Tr1 and Tr5 were completely inhibited at ten days of incubation on wheat straw by concentration of 3.6% (v/w) of the extract, and the strain of *P. ostreatus* at the same conditions showed a mycelial invasion delay of five days. It is suggested that the concentration of the fruiting body extract of *Pycnoporus* sp. at 3.6% could be effective for the control of green mold disease in mushroom cultures of the genus *Pleurotus*. It should be mentioned that the effect of the extract may also depend on the species, since each of them will have phenotypic and genotype characteristics that allow it to grow in adverse conditions. Marik et al. (2017) evaluated the aggressiveness of *T. aggressivum* (91 and 61%, respectively) and *T. pleuroti* (98 and 100%, respectively) against *A. bisporus* and *P. ostreatus*. In this work, *T. pleuroti* was the most resistant strain since the extract only decreased its growth by 38%. The extract of *Pycnoporus* sp. showed effectiveness, so it could be an alternative to control the green mold disease. It could be applied as a spray on the culture to prevent the invasion of *Trichoderma* spp., incorporating the extract during the incubation phase of *Pleurotus*, where the temperature increases due to the metabolic heat as a consequence of the mycelial growth of both fungi.

## Conclusions

The extract of the fruiting body of *Pycnoporus* sp. was able to reduce the growth of the strains of *T. pleuroti*, *T. atrobrunneum* and *T. aggressivum*, although the effect varied depending on the species. The *Pycnoporus* extract is suggested as an alternative for the control of green mold dis-

ease since the effect on the growth of *P. ostreatus* strain was minimal. The mushrooms are a source of obtaining metabolites that have antifungal activity, and they are a potential alternative to counteract the effects of *Trichoderma* in the cultivation of *Pleurotus*.

## Author contributions

ATO, MTT, PC and MLAU carried out field and laboratory work. MTT, GDG, EV and PC analyzed and interpreted the results. MTT and GDG wrote the manuscript. All authors read and approved the final manuscript.

## Funding

The National Council for Science and Technology (CONACYT) awarded a scholarship to Talavera-Ortiz (710481) for postgraduate studies of Master of Management of Natural Resources of the Center for Biological Research (Universidad Autónoma del Estado de Morelos).

## Acknowledgements

The authors thank CREGENHC (Centro de Recursos Genéticos de Hongos Comestibles, del Colegio de Postgraduados) for the donation of the *T. aggressivum* strain (CPM-113) and Vijai K. Gupta for reviewing the writing of the manuscript.

## Literature cited

- Al-Fatimi, M., G. Schröder, H. Kreisel and U. Lindequist. 2013. Biological activities of selected basidiomycetes from Yemen. Die Pharmazie-An International Journal of Pharmaceutical Sciences 68(3): 221-226.
- Beyer, D. M., P. J. Wuest and J. J. Kremser. 2000. Evaluation of epidemiological factors and mushroom substrate characteristics influencing the occurrence and development of *Trichoderma green* mold. In: Van Griensven, L. J. L. D. (ed.). Science and Cultivation of Edible Fungi. CRC Press. Maastricht, Netherlands. Pp. 633-640.
- Calonge, F. D. 2011. Hongos medicinales. Ed. Mundi-Prensa. Soria, España. 130 pp.
- Chaverri, P., F. Branco-Rocha, W. Jaklitsch, R. Gazis, T. Degenkolb and G. J. Samuels. 2015. Systematics of the *Trichoderma harzia-num* species complex and the re-identification of commercial biocontrol strains. Mycologia 107(3): 558-590. DOI: <https://doi.org/10.3852/14-147>

- Chen, X., C. P. Romaine, M. D. Ospina-Giraldo and D. J. Royse. 1999. A polymerase chain reaction-based test for the identification of *Trichoderma harzianum* biotypes 2 and 4, responsible for the worldwide green mold epidemic in cultivated *Agaricus bisporus*. *Applied Microbiology and Biotechnology* 52(2): 246-250. DOI: <https://doi.org/10.1007/s002530051516>
- Choi, I. Y., S. B. Hong and M. C. Yadav. 2003. Molecular and morphological characterization of green mold, *Trichoderma* spp. isolated from oyster mushrooms. *Mycobiology* 31(2): 74-80. DOI: <https://doi.org/10.4489/MYCO.2003.31.2.074>
- Danesh, Y. R., E. M. Goltapeh and H. Rohani. 2000. Identification of *Trichoderma* species causing green mold in button mushroom farms, distribution and their relative abundance. In: Van Griensven L. J. L. D. (ed.). *Science and Cultivation of Edible Fungi*. CRC Press. Maastricht, Netherland. Pp. 653-659.
- Díaz-Godínez, G., M. Téllez-Téllez, A. Rodríguez, V. Obregón-Barbosa, M. L. Acosta-Urdapilleta and E. Villegas. 2016. Enzymatic, antioxidant, antimicrobial, and insecticidal activities of *Pleurotus pulmonarius* and *Pycnoporus cinnabarinus* grown separately in an airlift reactor. *BioResources* 11(2): 4186-4200. DOI: <https://doi.org/10.15376/biores.11.2.4186-4200>
- Gea, F. J. 2009. First report of *Trichoderma pleurotum* on oyster mushroom crops in Spain. *Journal of Plant Pathology* 91: 504.
- Goltapeh, E. M. and Y. R. Danesh. 2000. Studies on interaction between *Trichoderma* species and *Agaricus bisporus* mycelium. In: Van Griensven, L. J. L. D. (ed.). *Science and Cultivation of Edible Fungi*. CRC Press. Maastricht, Netherlands. Pp. 661-666.
- Guzmán, G. 1994. Los hongos en la medicina tradicional de Mesoamérica y de México. *Revista Iberoamericana de Micología* 11(3): 81-85.
- Guzmán, G. 2003. Fungi in the Maya Culture: Past, Present and Future. In: Gómez-Pompa, A., M. F. Allen, S. L. Fedick and J. J. Jiménez-Osorio (eds.). *The Lowland Maya Area*. Food Products Press. Nueva York, USA. Pp. 315-325.
- Guzmán, G. 2008. Diversity and use of traditional Mexican medicinal fungi. A review. *International Journal of Medicinal Mushrooms* 10(3): 209-217. DOI: <https://doi.org/10.1615/IntJMedMushr.v10.i3.20>
- Hatvani, L., P. Sabolić, S. Kocsubé, L. Kredics, D. Cifra, C. Vágvölgyi, J. Kaliterna, D. Ivić and I. Kosalec. 2012. The first report on mushroom green mould disease in Croatia. *Arhiv za Higijenu Rada i Toksikologiju* 63(4): 481-486. DOI: <https://doi.org/10.2478/10004-1254-63-2012-2220>
- Hatvani, L., Z. Antal, L. Manczinger, A. Szekeres, I. S. Druzhinina, C. P. Kubicek, A. Nagy, E. Nagy, C. Vágvölgyi and L. Kredics. 2007. Green mould diseases of *Agaricus* and *Pleurotus* spp. are caused by related but phylogenetically different *Trichoderma* species. *Phytopathology* 97(4): 532-537. DOI: <https://doi.org/10.1094/PHYTO-97-4-0532>
- Index Fungorum. 2019. Index Fungorum database. <http://www.indexfungorum.org/Names/Names.asp> (consulted March, 2019).
- Kim, J. W., S. Kwon and H. J. Kang. 1995. Studies on the pathogenic *Pseudomonas* causing bacterial disease of cultivated mushroom in Korea. *Korean Journal Plant Pathology* 11: 353-360.
- Kim, K. Y., G. J. Lee, M. G. Ha, T. H. Lee and J. D. Lee. 2000. Intrageneric relationships of *Trichoderma* based on internal transcribed spacers and 5.8S rDNA nucleotide sequences. *Mycobiology* 28(1): 11-16. DOI: <https://doi.org/10.1080/12298093.2000.12015716>
- Komoń-Zelazowska, M., J. Bissett, D. Zafari, L. Hatvani, L. Manczinger, S. Woo, M. Lorito, L. Kredics, C. P. Kubicek and S. I. Druzhinina. 2007. Genetically closely related but phenotypically divergent *Trichoderma* species cause green mold disease in oyster mushroom farms worldwide. *Applied and Environmental Microbiology* 73(22): 7415-7426. DOI: <https://doi.org/10.1128/AEM.01059-07>
- Kredics, L., L. Hatvani, L. Antal, L. Manczinger, S. I. Druzhinina, P. C. Kubicek, A. Szekeres, A. Nagy, C. Vágvölgyi and E. Nagy. 2006. Green mold disease of oyster mushroom in Hungary and Transylvania. *Acta Microbiologica et Immunologica Hungarica* 53: 306-307.
- Kredics, L., S. Kocsubé, L. Nagy, M. Komoń-Zelazowska, L. Manczinger, E. Sajben, A. Nagy, C. Vágvölgyi, C. P. Kubicek, I. S. Druzhinina and L. Hatvani. 2009. Molecular identification of *Trichoderma* species associated with *Pleurotus ostreatus* and natural substrates of the oyster mushroom. *Microbiology Letters* 300(1): 58-67. DOI: <https://doi.org/10.1111/j.1574-6968.2009.01765.x>
- Marik, T., P. Urbán, C. Tyagi, A. Szekeres, B. Leitgeb, M. Vágvölgyi, L. Manczinger, I. S. Druzhinina, C. Vágvölgyi and L. Kredics. 2017. Diversity profile and dynamics of peptaibols produced by green mould *Trichoderma* species in interactions with their hosts *Agaricus bisporus* and *Pleurotus ostreatus*. *Chemistry and Biodiversity* 14(6): e1700033. DOI: <https://doi.org/10.1002/cbdv.201700033>

- Martínez-Alfaro, M. A., E. Pérez-Silva and E. Aguirre-Acosta. 1983. Et-nomicología y exploraciones micológicas en la Sierra Norte de Puebla. *Scientia Fungorum* 2(18): 51-63.
- NCBI. 2019. National Center for Biotechnology Information. <https://blast.ncbi.nlm.nih.gov> (consulted March, 2019).
- Oei, P. 1991. Manual on mushroom cultivation: techniques, species and opportunities for commercial application in developing countries. TOOL Publications. Wageningen, Netherlands. 249 pp.
- Park, M. S., K. S. Bae and S. H. Yu. 2004. Molecular and morphological analysis of *Trichoderma* isolates associated with green mold epidemic of oyster mushroom in Korea. *Journal of Huazhong Agricultural University* 23(1): 157-164.
- Park, M. S., K. S. Bae and S. H. Yu. 2006. Two new species of *Trichoderma* associated with green mold of oyster mushroom cultivation in Korea. *Mycobiology* 34(3): 111-113. DOI: <https://doi.org/10.4489/MYCO.2006.34.3.111>
- Qiu, Z., X. Wu, J. Zhang and C. Huang. 2017. High temperature enhances the ability of *Trichoderma asperellum* to infect *Pleurotus ostreatus* mycelia. *PloS ONE* 12(10): e0187055. DOI: <https://doi.org/10.1371/journal.pone.0187055>
- Reyes, Q. C. K. 2013. Propiedades antifúngicas diferenciadas de extractos vegetales como estrategia potencial para la prevención, manejo y control del moho verde (*Trichoderma* spp.): En el cultivo de los hongos comestibles (*Lentinula*, *Pleurotus*). Tesis de doctorado. Colegio de Posgraduados (COLPOS). Texcoco, México.
- Royse, D. J. 2014. A global perspective on the high five: *Agaricus*, *Pleurotus*, *Lentinula*, *Auricularia* & *Flammulina*. In: Singh, M. (ed.). *Proceedings of the 8th International Conference on Mushroom Biology and Mushroom Products (ICMBMP8)*. ICAR-Directorate of Mushroom Research. New Delhi, India. 639 pp.
- Samuels, G. L. and H. Prakash. 2015. *Trichoderma*: identification and agricultural applications (No. LC-0862). The American Phytopathological Society. Minnesota USA. 196 pp.
- Sharma, S. R. and B. Vijay. 1996. Yield loss in *Pleurotus ostreatus* spp. caused by *Trichoderma viride*. *Mushroom Research* 5: 19-22.
- Sobal, C. M., A. P. Morales, Q. M. Bonilla, S. W. Martínez and D. Martínez-Carrera. 2016. Biotecnología, innovación y desarrollo en la cadena de valor con base en los recursos genéticos de los hongos comestibles, medicinales y funcionales. In: Martínez-Carrera, D. and J. J. Ramírez (eds.). *Ciencia, Tecnología e Innovación en el Sistema Agroalimentario de México*. Colegio de Postgraduados-Academia Mexicana de Ciencias-Consejo Nacional de Ciencia y Tecnología-Universidad Popular Autónoma del Estado de Puebla-Instituto de Micología Neotropical Aplicada Press. Texcoco, México. Pp. 761-779.
- Sobieralski, K., M. Siwulski, M. Kommon-Żelazowska, L. Błaszczuk, I. Sas-Golak and D. Frużyńska-Jóźwiak. 2012. Impact of *Trichoderma pleurotum* and *T. pleuroticola* isolates on yielding of *Pleurotus ostreatus* (FR.) Kumm. *Journal of Plant Protection Research* 52(1): 165-168. DOI: <https://doi.org/10.2478/v10045-012-0025-4>
- Soković, M. and L. J. van Griensven. 2006. Antimicrobial activity of essential oils and their components against the three major pathogens of the cultivated button mushroom, *Agaricus bisporus*. *European Journal of Plant Pathology* 116(3): 211-224. DOI: <https://doi.org/10.1007/s10658-006-9053-0>
- StatSoft Inc. 2004. STATISTICA (data analysis software system) version 7.0. Tulsa, USA. <https://statistica.software.informer.com/7.0/> (consulted March, 2019).
- Téllez-Téllez, M., G. Díaz-Godínez and C. Sánchez. 2003. Physiology of colony *Pleurotus pulmonarius* grown on media overlaid with a cellophane membrane. *Applied Microbiology and Biotechnology* 63(2): 212-216. DOI: <https://doi.org/10.1007/s00253-003-1387-3>
- Teoh, Y. P., M. M. Don and S. Ujang. 2011. Media selection for mycelia growth, antifungal activity against wood-degrading fungi, and GC-MS study by *Pycnoporus sanguineus*. *BioResources* 6(3): 2719-2731. DOI: <https://doi.org/10.15376/biores.6.3.2719-2731>
- Wang, G., X. Cao, X. Ma, M. Guo, C. Liu, L. Yan and Y. Bian. 2016. Diversity and effect of *Trichoderma* spp. associated with green mold disease on *Lentinula edodes* in China. *Microbiology Open* 5(4): 709-718. DOI: <https://doi.org/10.1002/mbo3.364>
- Woo, S. L., P. Di Benedetto, M. Senatore, K. Abadi, S. Gigante, I. Soriente, S. Ferraioli, F. Scala and M. Lorito. 2004. Identification and characterization of *Trichoderma* species aggressive to *Pleurotus* in Italy. *Journal of Zhejiang University Agriculture and Life Sciences* 30: 469-470.
- Zehr, E. I. 1978. Methods for evaluating plant fungicides, nematocides, and bactericides. Prepared jointly by the American Phytopathological Society and the Society of Nematologists. American Phytopathological Society. St. Paul, Minnesota, USA. 141 pp.